

# A Cauchy-Schwarz Type Inequality for Bilinear Integrals on Positive Measures

Nils Ackermann

April 2004

## Abstract

If  $W: \mathbb{R}^n \rightarrow [0, \infty]$  is Borel measurable, define for  $\sigma$ -finite positive Borel measures  $\mu, \nu$  on  $\mathbb{R}^n$  the bilinear integral expression

$$I(W; \mu, \nu) := \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} W(x - y) d\mu(x) d\nu(y) .$$

We give conditions on  $W$  such that there is a constant  $C \geq 0$ , independent of  $\mu$  and  $\nu$ , with

$$I(W; \mu, \nu) \leq C \sqrt{I(W; \mu, \mu) I(W; \nu, \nu)} .$$

Our results apply to a much larger class of functions  $W$  than known before.

## 1. Introduction and Results

Given a Borel function  $W: \mathbb{R}^n \rightarrow [0, \infty]$ , for  $\sigma$ -finite positive measures  $\mu, \nu$  on  $\mathbb{R}^n$  define the integral

$$I(W; \mu, \nu) := \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} W(x - y) d\mu(x) d\nu(y) .$$

Denote for  $C \geq 0$  by  $\mathcal{W}(n, C)$  the class of Borel functions  $W: \mathbb{R}^n \rightarrow [0, \infty]$  such that for all  $\sigma$ -finite positive measures  $\mu, \nu$  on  $\mathbb{R}^n$

$$(1.1) \quad I(W; \mu, \nu) \leq C \sqrt{I(W; \mu, \mu) I(W; \nu, \nu)}$$

holds. Moreover, denote

$$\mathcal{W}(n) := \bigcup_{C \geq 0} \mathcal{W}(n, C) .$$

If  $W$  is an even function and the symmetric bilinear form  $I(W; \cdot, \cdot)$  is positive semidefinite, then  $W \in \mathcal{W}(n, 1)$  (Cauchy-Schwarz' inequality). Hence we may regard (1.1) as a generalized form of the Cauchy-Schwarz inequality.

An even function  $W$  such that  $I(W; \cdot, \cdot)$  is positive semidefinite is called *positive definite*. Roughly speaking, positive definiteness of a function corresponds to non negativity of its Fourier transform [5, 6]. The only result regarding (1.1) the author is aware of that goes beyond positive definite functions is given by Mattner [4, Sect. 5.1]: If  $\|\cdot\|$  is any norm on  $\mathbb{R}^n$ ,  $h: [0, \infty) \rightarrow [0, \infty]$  is decreasing, and  $W$  is given by  $W(x) := h(\|x\|)$ , then  $W \in \mathcal{W}(n)$ . Theorem 1.5 below recovers this statement and extends it by allowing  $h$  to be non monotone. Theorem 1.2, the main result of the present paper, yields a criterion for membership in  $\mathcal{W}(n)$  for functions  $W$  that can not be written as  $h \circ p$  with a seminorm  $p$  on  $\mathbb{R}^n$ .

The study of property (1.1) is motivated by the partial differential equation

$$(1.2) \quad -\Delta u + Vu = (W * u^2)u \quad u \in H^1(\mathbb{R}^n).$$

Here  $*$  denotes convolution,  $V$  in  $L^\infty(\mathbb{R}^n)$  is periodic, and 0 lies in a gap of the spectrum of  $(-\Delta + V)$ , cf. [1]. One is interested in the existence of nontrivial solutions to (1.2). For the special case  $n = 3$  and  $W(x) = 1/\|x\|_2$  the problem was settled in [2] by using the fact that this particular function  $W$  is positive definite. In [1] it is shown that  $W \in \mathcal{W}(n)$  (together with appropriate growth conditions) is sufficient to obtain a nontrivial solution of (1.2).

## 1.1. Main Results

The statement of our Theorems requires to introduce some notation and definitions. For a topological space  $X$  denote by  $\mathcal{P}(X)$  the set of Borel functions  $f: X \rightarrow [0, \infty]$ . For  $n \in \mathbb{N}$  denote by  $\mathcal{C}(n)$  the class of subsets of  $\mathbb{R}^n$  that are closed, convex, and symmetric (i.e.  $-A = A$ ). The dimension  $\dim A$  of a convex subset  $A$  of  $\mathbb{R}^n$  is the dimension of the affine hull of  $A$ .

**Definition 1.1.** For  $X, A \subseteq \mathbb{R}^n$ ,  $X \neq \emptyset$ , put

$$\kappa(X, A) := \inf\{m \in \mathbb{N} \mid X \text{ can be covered by } m \text{ translates of } A\}$$

and

$$\alpha(X) := \inf\{m \in \mathbb{N} \mid \exists A \in \mathcal{C}(n): \dim A = n, A \subseteq X \text{ and } \kappa(X, A) = m\}.$$

For  $X = \emptyset$  set  $\kappa(\emptyset, A) := 0$  and  $\alpha(\emptyset) := 0$ .

Given a set  $X$ , a map  $W: X \rightarrow \mathbb{R}$  and  $t$  in  $\mathbb{R}$  denote

$$[W]_t := \{x \in X \mid W(x) \geq t\}.$$

Furthermore, define the class  $\mathcal{A}(n)$  by

$$\mathcal{A}(n) := \left\{ W \in \mathcal{P}(\mathbb{R}^n) \mid \limsup_{t \rightarrow 0} \alpha([W]_t) + \limsup_{t \rightarrow \infty} \alpha([W]_t) < \infty \right\}.$$

Our main result then reads:

**Theorem 1.2.** For every  $n$  in  $\mathbb{N}$  the inclusion  $\text{conv}(\mathcal{A}(n)) \subseteq \mathcal{W}(n)$  holds.

**Remark 1.3.** It will be shown in the proof of Theorem 1.2 that  $\mathcal{W}(n)$  is a convex cone. Obviously,  $\mathcal{A}(n)$  is a cone. The Example 1.6 given below demonstrates that  $\mathcal{A}(n)$  is not convex.

The present author does not know whether a function in  $\mathcal{W}(n)$  that is sufficiently regular, say continuous, must necessarily belong to  $\text{conv}(\mathcal{A}(n))$ .

A simpler criterion for membership in  $\mathcal{W}(n)$  can be formulated in the case of the composition of a map with a seminorm. To state it we introduce further concepts and notation.

**Definition 1.4.** For a subset  $Y$  of  $[0, \infty)$  put  $\lambda(Y) := \sup\{t > 0 \mid [0, t] \subseteq Y\}$  and

$$\beta(Y) := \begin{cases} 0 & Y = \emptyset \\ \infty & \lambda(Y) = -\infty \text{ and } Y \neq \emptyset \\ \sup(Y)/\lambda(Y) & \text{otherwise.} \end{cases}$$

Here we set  $\infty/a := \infty$  if  $a > 0$  and  $\infty/\infty := 1$ .

We introduce

$$\mathcal{B} := \left\{ h \in \mathcal{P}([0, \infty)) \mid \limsup_{t \rightarrow 0} \beta([h]_t) + \limsup_{t \rightarrow \infty} \beta([h]_t) < \infty \right\}.$$

Our second result then reads:

**Theorem 1.5.** Suppose that  $h \in \mathcal{P}([0, \infty))$  and that  $p$  is a seminorm on  $\mathbb{R}^n$ . If  $h \in \mathcal{B}$  then  $h \circ p \in \mathcal{A}(n)$ . If  $h \circ p \in \mathcal{A}(n)$  and  $\text{codim}(\ker p) \geq 2$  then  $h \in \mathcal{B}$ .

We provide some examples to illustrate the concepts introduced so far:

**Example 1.6.** Denote by  $h$  the characteristic function of  $[0, 1]$ , taken as a map from  $[0, \infty)$  into  $[0, \infty]$ . Then  $h \in \mathcal{B}$ . For  $i = 1, 2$  define  $W_i$  as a map in  $\mathcal{P}(\mathbb{R}^2)$  by  $W_i(x_1, x_2) := h(|x_i|)$ . Theorem 1.5 implies that  $W_i \in \mathcal{A}(2)$  for  $i = 1, 2$ , but clearly  $W := W_1 + W_2 \notin \mathcal{A}(2)$ . Since  $\mathcal{A}(2)$  is a cone this implies that  $\mathcal{A}(2)$  is not convex. Nevertheless,  $W \in \mathcal{W}(2)$  by Theorem 1.2 and since  $\mathcal{W}(2)$  is a convex cone.

**Example 1.7.** We construct a function  $W$  in  $\mathcal{A}(n)$  that is not even, and hence is neither positive definite nor of the form  $h \circ p$  with  $h$  in  $\mathcal{P}([0, \infty))$  and  $p$  a seminorm on  $\mathbb{R}^n$ . Pick  $x_0$  in  $\mathbb{R}^n \setminus \{0\}$  and set

$$W_0(x) := \frac{1}{\|x\|_2}$$

$$W(x) := W_0(x) + W_0(x - x_0).$$

Denoting by  $D(r, x)$  the closed ball of radius  $r > 0$  with center  $x$ , it follows easily that

$$D(1/t, 0) \subseteq [W]_t \subseteq D(2/t, 0) \cup D(2/t, x_0)$$

for all  $t > 0$ . This implies that  $W \in \mathcal{A}(n)$ .

**Example 1.8.** We show that the assumption on  $\text{codim}(\ker p)$  used in Theorem 1.5 is not purely technical. If  $p$  is a seminorm on  $\mathbb{R}^n$  with  $\text{codim}(\ker p) = 0$  then trivially  $h \circ p \in \mathcal{A}(n)$  for arbitrary  $h$  in  $\mathcal{P}([0, \infty))$ . Given the seminorm  $p(x) := |x|$  in  $\mathbb{R}$  with  $\text{codim}(\ker p) = 1$ , we construct  $h$  in  $\mathcal{P}([0, \infty))$  such that  $W := h \circ p \in \mathcal{A}(1)$  but  $h \notin \mathcal{B}$ . Put

$$h(s) := \begin{cases} \infty & s = 0 \\ \exp(-(k-1)^2) & s = \exp(k^2) \text{ for some } k \text{ in } \mathbb{N} \\ 1/s & \text{otherwise.} \end{cases}$$

For  $t > 1$  we obtain  $[h]_t = [0, 1/t]$ , and for  $0 < t \leq 1$  we obtain

$$(1.3) \quad [h]_t = [0, 1/t] \cup \left\{ \exp\left(\left(1 + \left[\sqrt{-\log t}\right]\right)^2\right) \right\}.$$

Recall that  $[a]$  denotes the largest integer less than or equal to  $a$  if  $a \in \mathbb{R}$ . From (1.3) it is clear that  $\alpha([W]_t) \leq 3$  for all  $t \geq 0$ , so  $W \in \mathcal{A}(1)$ . On the other hand, for  $t_k := \exp(-k^2)$  we find

$$\beta([h]_{t_k}) = \exp((1+k)^2) \exp(-k^2) = \exp(1+2k)$$

and therefore  $\limsup_{t \rightarrow 0} \beta([h]_t) = \infty$ . Hence  $h \notin \mathcal{B}$ .

## 1.2. General Notation

In  $\mathbb{R}^n$  denote by  $\|\cdot\|_p$  for  $p$  in  $[1, \infty]$  the standard  $l^p(n)$ -norm. In the case of  $p = 2$  we write  $x \cdot y$  for the standard Euclidean scalar product of elements  $x, y$  in  $\mathbb{R}^n$ . If  $V$  is a subspace of  $\mathbb{R}^n$ , denote by  $V^\perp$  the orthogonal subspace with respect to the standard scalar product.

The power set of a set  $X$  will be written  $2^X$ . The cardinality of  $X$  is denoted by  $|X|$ . Some operators used are:  $\text{conv } A$  for the convex hull of  $A$ ,  $\text{cl } A$ ,  $\text{int } A$ , and  $\partial A$  for closure, interior, and boundary of a subset  $A$  of a topological space.

A parallelotope is a rectangular parallelepiped.

## 2. Some Convex Geometry

The next Lemma allows us to deal with unbounded sets in  $\mathcal{C}(n)$  in a convenient manner.

**Lemma 2.1.** *If  $A \in \mathcal{C}(n)$  then there is a unique subspace  $V$  of  $\mathbb{R}^n$  such that  $B := A \cap V^\perp \in \mathcal{C}(n)$  is compact and  $A = B + V$ .*

*Proof.* First we remark: If a set  $A$  in  $\mathcal{C}(n)$  includes a ray (a set  $\{x + ty \mid t \geq 0\}$  for some  $x, y$  in  $\mathbb{R}^n$ ), then it includes the 1-dimensional subspace parallel to that ray. If  $A$  includes a translate of a subspace  $V$  of  $\mathbb{R}^n$  then  $A$  includes  $V$ .

Now fix  $A$  in  $\mathcal{C}(n)$ . From [3, Lemma 2.5.4] we obtain a unique subspace  $V$  of  $\mathbb{R}^n$  of maximal dimension such that a translate of  $V$  and thus  $V$  is included in  $A$ . Moreover, by that lemma it also holds that  $B := A \cap V^\perp \in \mathcal{C}(n)$  does not include a line (the translate of a 1-dimensional subspace) and  $A = B + V$ . If  $B$  was not bounded then it included a ray

by [3, Lemma 2.5.1]. Since  $B$  is symmetric it therefore included a line also. Contradiction. Since  $A$  is closed  $B$  must therefore be compact.

If, on the other hand, for some subspace  $V$  of  $\mathbb{R}^n$ ,  $B = A \cap V^\perp$  is compact and  $A = B + V$ , then  $V$  is included in  $A$ . If  $A$  includes a translate of another subspace  $W$ , and thus includes  $W$ , then  $W \subseteq V$ . Hence  $V$  has maximal dimension among the subspaces included in  $A$ , and it is unique, again by Lemma 2.5.4 *loc.cit.*  $\square$

**Definition 2.2.** We call the pair  $(B, V)$  given for  $A$  in  $\mathcal{C}(n)$  by Lemma 2.1 the *splitting of  $A$* .

**Definition 2.3.** Denote for  $X \subseteq \mathbb{R}^n$  by

$$\text{ccs } X := \text{cl}(\text{conv } \frac{1}{2}(X - X)) \in \mathcal{C}(n)$$

the closed convex hull of the symmetrization of  $X$ .

**Remark 2.4.** For  $A, B \subseteq \mathbb{R}^n$  we have  $\text{conv}(A + B) = \text{conv } A + \text{conv } B$ . Thus

$$\text{ccs } X = \text{cl } \frac{1}{2}(\text{conv } X - \text{conv } X) .$$

From this also follows that  $\text{ccs}(X + Y) = \text{ccs } X + \text{ccs } Y$  if one of  $X$  and  $Y$  is relative compact. Moreover,  $\text{ccs } A = A$  if  $A \in \mathcal{C}(n)$ .

**Definition 2.5.** If  $X \subseteq \mathbb{R}^n$  and  $(A, V)$  is the splitting of  $\text{ccs } X$ , put  $\gamma(X) := \dim V$ .

**Lemma 2.6.** *The map  $\gamma: 2^{\mathbb{R}^n} \rightarrow \{0, 1, 2, \dots, n\}$  is monotone increasing with respect to the partial order induced on  $2^{\mathbb{R}^n}$  by inclusion. If  $X \subseteq Y \subseteq \mathbb{R}^n$  and  $\gamma(X) = \gamma(Y)$ , then from  $A \in \mathcal{C}(n)$  with  $\dim A = n$  and  $\kappa(X, A) < \infty$  it follows that  $\kappa(Y, A) < \infty$ .*

*Proof.* Monotonicity of  $\gamma$  is obvious. Fix  $X \subseteq Y$  with  $\gamma(X) = \gamma(Y)$ , and suppose we are given  $A$  in  $\mathcal{C}(n)$  with  $\dim A = n$  and  $\kappa(X, A) < \infty$ . Let  $(B, V)$  be the splitting of  $A$  and let  $\mathcal{I} \subseteq \mathbb{R}^n$  be finite with  $X \subseteq \mathcal{I} + A = \mathcal{I} + B + V$ . Since  $\mathcal{I} + B$  is compact, in view of Remark 2.4 we obtain

$$(2.1) \quad \text{ccs } X \subseteq \text{ccs}(\mathcal{I} + B + V) = \text{ccs}(\mathcal{I} + B) + V .$$

Since  $\text{ccs } X \subseteq \text{ccs } Y$  and  $\gamma(X) = \gamma(Y)$  there is a subspace  $W$  of  $\mathbb{R}^n$  with  $\dim W = \gamma(X)$  and there are splittings  $(B_1, W)$  and  $(B_2, W)$  of  $\text{ccs } X$  and  $\text{ccs } Y$  respectively, with  $B_1 \subseteq B_2$ . Put  $A_1 := A \cap W^\perp$ . Now (2.1) implies  $W \subseteq V$ , and hence  $A_1 + W = A$ . Therefore  $\dim A = n$  yields  $\dim A_1 = \dim W^\perp = n - \gamma(X)$ , and  $\text{relint } A_1$  (the interior of  $A_1$  relative to the smallest subspace including  $A_1$ ) is open in  $W^\perp$ . Since  $B_2 \subseteq W^\perp$  is compact there is a finite set  $\mathcal{J} \subseteq W^\perp$  with  $B_2 \subseteq \mathcal{J} + A_1$ . It follows that

$$Y \subseteq \text{ccs } Y = B_2 + W \subseteq \mathcal{J} + A_1 + W = \mathcal{J} + A$$

and thus  $\kappa(Y, A) < \infty$ .  $\square$

**Lemma 2.7.** *For all  $n$  in  $\mathbb{N}$  there is a constant  $C_1(n) \geq 0$  such that for all  $A$  in  $\mathcal{C}(n)$  with  $\dim A = n$  the following hold:*

a)  $\kappa(A, \frac{1}{2}A) \leq C_1(n)$

b) *there is a discrete subgroup  $G$  of the additive group of  $\mathbb{R}^n$  such that  $\mathbb{R}^n = G + A$  and  $\sup_{x \in \mathbb{R}^n} |(x + 3A) \cap G| \leq C_1(n)$ .*

*Proof.* From [7, Lemma 2.4] we obtain for all  $m$  in  $\mathbb{N}$  a constant  $C_2(m)$ , monotone increasing in  $m$ , such that for every  $m$ -dimensional compact  $B$  in  $\mathcal{C}(m)$  there is a parallelotope  $P \subseteq \mathbb{R}^m$ , centered at the origin, with

$$(2.2) \quad P \subseteq B \subseteq C_2(m)P .$$

Now set

$$C_1(n) := [3C_2(n) + 1]^n$$

where  $[a]$  denotes the largest integer below or equal to  $a$  if  $a \in \mathbb{R}$ .

Fix  $A$  in  $\mathcal{C}(n)$  and let  $(B, V)$  be the splitting of  $A$ . Since  $\dim A = n$  we have  $\dim B + \dim V = n$ . We may assume  $\dim B = m$  and  $V = \{0\} \times \mathbb{R}^{n-m}$  as a subspace of  $\mathbb{R}^n$ . We identify  $\mathbb{R}^m$  with  $\mathbb{R}^m \times \{0\} \subseteq \mathbb{R}^n$  so that  $B \subseteq \mathbb{R}^m$ , and we choose a parallelotope  $P \subseteq \mathbb{R}^m$  for  $B$  as in (2.2). Then from  $2C_2(m) \leq 3C_2(n)$  and the definition of  $C_1(n)$  we obtain

$$\kappa(A, \frac{1}{2}A) = \kappa(B, \frac{1}{2}B) \leq \kappa(C_2(m)P, \frac{1}{2}P) \leq \kappa(3C_2(n)P, P) \leq C_1(n) .$$

For the second assertion we use for  $P$  from above the representation

$$P = [-r_1, r_1] \times [-r_2, r_2] \times \cdots \times [-r_m, r_m]$$

with some  $r_1, r_2, \dots, r_m > 0$  and put  $G_0 := 2r_1\mathbb{Z} \times 2r_2\mathbb{Z} \times \cdots \times 2r_m\mathbb{Z} \subseteq \mathbb{R}^m$ . Then  $G_0$  is an additive subgroup of  $\mathbb{R}^m$  with  $G_0 + B \supseteq G_0 + P = \mathbb{R}^m$ . Now set  $G := G_0 \times \{0\} \subseteq \mathbb{R}^n$ . Then  $G + A = G + B + V = \mathbb{R}^n$ . On the other hand we have for every  $x$  in  $\mathbb{R}^n$

$$(x + 3A) \cap G = (x + 3B) \cap G \subseteq (x + 3C_2(m)P) \cap G \subseteq (x + 3C_2(n)P) \cap G$$

and hence

$$|(x + 3A) \cap G| \leq |(x + 3C_2(n)P) \cap G| \leq C_1(n) .$$

This completes the proof. □

**Lemma 2.8.** *Suppose that  $p$  is a seminorm on  $\mathbb{R}^n$  and that  $Y \subseteq [0, \infty)$ . Put  $X := p^{-1}(Y)$ . Then  $\alpha(X) \leq C_3(n)\beta(Y)^n$  for some constant  $C_3(n)$ . If  $\text{codim}(\ker p) \geq 2$ , then  $\alpha(X) \geq \beta(Y)/2$ .*

*Proof.* For  $r > 0$  put  $A(r) := \{x \in \mathbb{R}^n \mid p(x) \leq r\} \in \mathcal{C}(n)$ . Let  $(B(1), V)$  be the splitting of  $A(1)$  and put  $B(r) := rB(1)$  for  $r > 0$ . Then  $(B(r), V)$  is the splitting of  $A(r)$ . Moreover,  $V = \ker p$ . Set  $m := \text{codim } V$ , so  $\dim B(1) = m$ .

Define  $f, g: [0, \infty) \rightarrow \mathbb{N}$  by setting  $f(0) := g(0) := 1$  and, for  $t > 0$ ,  $f(t) := \kappa(\partial A(t), A(1)) = \kappa(\partial B(t), B(1))$  and  $g(t) := \kappa(A(t), A(1)) = \kappa(B(t), B(1))$ . Then  $f$  and  $g$  are monotone increasing,  $f \leq g$ , and

$$\begin{aligned} \kappa(\partial A(r), A(s)) &= f(r/s) \\ \kappa(A(r), A(s)) &= g(r/s) \end{aligned}$$

for  $r, s > 0$ . As in the beginning of the proof of Lemma 2.7 we obtain

$$(2.3) \quad g(t) = \kappa(B(t), B(1)) \leq \kappa(tC_2(m)P, P) = [tC_2(m) + 1]^m .$$

Here  $P \subseteq B(1)$  is a parallelotope chosen as for (2.2). If  $m \geq 2$  then

$$(2.4) \quad f(t) = \kappa(\partial B(t), B(1)) \geq t .$$

This can be seen as follows: Consider  $B(1)$  as a subset of  $\mathbb{R}^m$ . Fix  $x_0$  in  $\partial B(1)$  such that  $2\|x_0\|_2 = \text{diam } B(1)$ . Let  $Q$  be the orthogonal projection in  $\mathbb{R}^m$  onto  $\text{span}\{x_0\}$  and  $L := \ker Q$ . Then  $\dim L \geq 1$ . It follows that for every  $x$  in  $[-tx_0, tx_0]$  (the segment joining  $-tx_0$  and  $tx_0$ ) the set  $(x + L) \cap \partial B(t)$  is not empty. Moreover, from  $B(1) \in \mathcal{C}(n)$  it follows that  $x_0 + L$  is a supporting hyperplane for  $B(1)$ . If  $x_1, x_2, \dots, x_k \in \mathbb{R}^m$  are such that

$$\partial B(t) \subseteq \bigcup_{l=1}^k (x_l + B(1)) ,$$

from the above it is clear that then

$$[-tx_0, tx_0] \subseteq \bigcup_{l=1}^k (Qx_l + B(1))$$

and therefore  $k \geq [t + 1] \geq t$ . This yields (2.4).

Let us consider the case  $0 < \lambda(Y) \leq \sup Y < \infty$ . There is

$$\varepsilon \in [0, \lambda(Y)/2]$$

such that  $[0, \lambda(Y) - \varepsilon] \subseteq Y$ . It follows that

$$A(\lambda(Y) - \varepsilon) \subseteq X \subseteq A(\sup Y) .$$

Thus, using (2.3), we obtain

$$\alpha(X) \leq \kappa(A(\sup Y), A(\lambda(Y) - \varepsilon)) = g\left(\frac{\sup Y}{\lambda(Y) - \varepsilon}\right) \leq g(2\beta(Y)) \leq C_3(n)\beta(Y)^n$$

for some constant  $C_3(n) \geq 1$ .

There is  $\varepsilon$  in  $[0, \sup Y/2]$  such that  $\sup Y - \varepsilon \in Y$  and therefore

$$(2.5) \quad \partial A(\sup Y - \varepsilon) \subseteq X .$$

Every  $A$  in  $\mathcal{C}(n)$  with  $A \subseteq X$  is path connected, and satisfies  $0 \in A$ . Since  $p$  is continuous,  $p(A)$  is included in the path component of  $Y$  containing 0. Therefore  $p(A) \subseteq [0, \lambda(Y)]$  and  $A \subseteq A(\lambda(Y))$ . This shows that

$$\kappa(X, A) \geq \kappa(X, A(\lambda(Y)))$$

for all  $A$  in  $\mathcal{C}(n)$ . Hence we find for  $m \geq 2$ , applying (2.4) and (2.5):

$$\alpha(X) \geq \kappa(\partial A(\sup Y - \varepsilon), A(\lambda(Y))) = f\left(\frac{\sup Y - \varepsilon}{\lambda(Y)}\right) \geq f(\beta(Y)/2) \geq \beta(Y)/2 .$$

The case  $\lambda(Y) > 0, \sup(Y) = \infty$  is handled similarly, and in all other cases the assertion is trivial.  $\square$

### 3. Proof of the Theorems

Let us first prove Theorem 1.5. Suppose that we are given  $h \in \mathcal{P}([0, \infty))$  and a seminorm  $p$  on  $\mathbb{R}^n$ . Set  $W := h \circ p$ . Then  $[W]_t = p^{-1}([h]_t)$  for every  $t > 0$ . Now Lemma 2.8 yields  $\alpha([W]_t) \leq C\beta([h]_t)^n$  with some positive constant  $C$ . Moreover, if  $\text{codim}(\ker p) \geq 2$  Lemma 2.8 implies that  $\beta([h]_t) \leq 2\alpha([W]_t)$ . From these facts the theorem follows.

The proof of Theorem 1.2, taken up next, is divided into the following steps:

- (i)  $\mathcal{W}(n, C)$  is closed under increasing pointwise limits for every  $C \geq 0$ .
- (ii)  $\mathcal{W}(n, C)$  is a convex cone for every  $C \geq 0$ .

Now suppose that  $W \in \mathcal{P}(\mathbb{R}^n)$ .

- (iii) If  $A$  in  $\mathcal{C}(n)$  has dimension  $n$ , if  $\kappa(\text{supp } W, A) < \infty$ , if there is  $a > 0$  such that  $W \geq a$  on  $2A$ , and if  $W$  is bounded with  $b := \sup W(\mathbb{R}^n)$ , then  $W \in \mathcal{W}(n, C)$  for  $C := C_1(n)^3 \kappa(\text{supp } W, A) b/a$ , where  $C_1(n)$  is the constant given in Lemma 2.7.
- (iv) If  $\sup_{t \geq 0} \alpha([W]_t) < \infty$  then  $W \in \mathcal{W}(n, C)$  for some  $C \geq 0$ .
- (v) If  $\limsup_{t \rightarrow 0} \alpha([W]_t) + \limsup_{t \rightarrow \infty} \alpha([W]_t) < \infty$  then  $W \in \mathcal{W}(n, C)$  for some  $C \geq 0$ .

Theorem 1.2 is then a consequence of (ii) and (v).

Statements (i) and (ii) were proven in [4, Sect. 5.1]. For completeness we repeat the argument here. Suppose that  $C \geq 0$ . Fix two  $\sigma$ -finite positive Borel measures  $\mu, \nu$  on  $\mathbb{R}^n$ . If  $W$  is the pointwise limit of an increasing sequence of functions in  $\mathcal{W}(n, C)$ , then (1.1) follows from Lebesgue's Monotone Convergence Theorem. This proves (i) since  $\mu, \nu$  were chosen arbitrarily.

Consider the implication

$$(3.1) \quad \left( u \leq C\sqrt{vw} \quad \text{and} \quad x \leq C\sqrt{yz} \right) \quad \Rightarrow \quad (u+x)^2 \leq C^2(v+y)(w+z)$$

for  $u, v, w, x, y, z$  in  $[0, \infty)$ , which is a consequence of  $2\sqrt{vwyz} \leq vz + yw$ . If  $W_1, W_2 \in \mathcal{W}(n, C)$  then (3.1) implies that  $W_1 + W_2 \in \mathcal{W}(n, C)$ . Since  $\mathcal{W}(n, C)$  is a cone,  $\mathcal{W}(n, C)$  is convex.

To show (iii) choose a discrete additive subgroup  $G$  of  $\mathbb{R}^n$  for  $A$  as in Lemma 2.7b). Let  $\mathcal{I}$  be a finite subset of  $\mathbb{R}^n$  with  $\text{supp } W \subseteq \mathcal{I} + A$  and  $|\mathcal{I}| = \kappa(\text{supp } W, A)$ . Put  $\mathcal{J} := (\mathcal{I} + 3A) \cap G$ . From the choice of  $G$  it follows that

$$(3.2) \quad |\mathcal{J}| \leq C_1(n)|\mathcal{I}|.$$

Define  $\overline{W}: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  by  $\overline{W}(x, y) := W(x - y)$ . Then  $\overline{W}$  is a Borel function. We claim that

$$\text{supp } \overline{W} \subseteq \bigcup_{\substack{u, v \in G \\ u-v \in \mathcal{J}}} (u + A) \times (v + A).$$



To see this, suppose that  $(x, y) \in \text{supp } \overline{W}$ , or equivalently  $x - y \in \text{supp } W$ . There is  $w$  in  $\mathcal{I}$  such that  $x - y \in w + A$ , and there are  $u, v$  in  $G$  such that  $x \in u + A$  and  $y \in v + A$ . It follows that  $u - v \in x - y + 2A \subseteq w + 3A \subseteq \mathcal{I} + 3A$ . Also  $u - v \in G$  because  $G$  is a subgroup. This proves the claim.

Now Cauchy-Schwarz' inequality for sums yields

$$(3.3) \quad I(W; \mu, \nu) = \int_{\mathbb{R}^n \times \mathbb{R}^n} \overline{W} d(\mu \times \nu) \leq b \int_{\text{supp } \overline{W}} d(\mu \times \nu) \\ \leq b \sum_{\substack{u, v \in G \\ u - v \in \mathcal{J}}} \mu(u + A) \nu(v + A) \leq b \left( \sum_{\substack{u, v \in G \\ u - v \in \mathcal{J}}} \mu(u + A)^2 \sum_{\substack{u, v \in G \\ u - v \in \mathcal{J}}} \nu(v + A)^2 \right)^{\frac{1}{2}}.$$

We need to estimate the sums in the last term. For every  $x$  in  $\mathbb{R}^n$ , from  $A \in \mathcal{C}(n)$  it follows that the statement  $(u \in G \text{ and } x \in u + A)$  is equivalent to the statement  $u \in (x + A) \cap G$ . By the choice of  $G$  this leads to

$$|\{u \in G \mid x \in u + A\}| = |(x + A) \cap G| \leq |(x + 3A) \cap G| \leq C_1(n)$$

and thus for all  $x, y$  in  $\mathbb{R}^n$

$$(3.4) \quad |\{u \in G \mid (x, y) \in (u + A) \times (u + A)\}| \leq C_1(n)^2.$$

Also we have

$$(3.5) \quad \bigcup_{u \in G} (u + A) \times (u + A) \subseteq \{(x, y) \in \mathbb{R}^n \times \mathbb{R}^n \mid x - y \in 2A\} =: D$$

and  $\overline{W} \geq a$  on  $D$ . Using (3.2), (3.4) and (3.5) we calculate

$$\sum_{\substack{u, v \in G \\ u - v \in \mathcal{J}}} \mu(u + A)^2 = |\mathcal{J}| \sum_{u \in G} \mu(u + A)^2 = |\mathcal{J}| \sum_{u \in G} \int_{(u+A) \times (u+A)} d(\mu \times \mu) \\ \leq C_1(n)^2 |\mathcal{J}| \int_D d(\mu \times \mu) \leq \frac{C_1(n)^3 |\mathcal{I}|}{a} \int_{\mathbb{R}^n \times \mathbb{R}^n} \overline{W} d(\mu \times \mu) \\ = \frac{C_1(n)^3 |\mathcal{I}|}{a} I(W; \mu, \mu),$$

a similar estimate holding for the sum over  $\nu(v + A)^2$ . This proves (iii) in view of (3.3).

To show (iv) suppose that  $M := \sup_{t \geq 0} \alpha([W]_t) < \infty$ . For  $m$  in  $\mathbb{N}$  and  $1 \leq i \leq m2^m$  define  $W_{m,i}$  and  $W_m$  in  $\mathcal{P}(\mathbb{R}^n)$  by setting

$$W_{m,i} := \frac{1}{2^m} \chi_{[W]_{i/2^m}} \\ W_m := \sum_{i=1}^{m2^m} W_{m,i}.$$

Here  $\chi_A$  denotes for  $A \subseteq \mathbb{R}^n$  the characteristic function of  $A$ . The sequence  $(W_m)$  is increasing and converges pointwise to  $W$ . Fix  $m$  and  $i$ . There is  $A$  in  $\mathcal{C}(n)$  such that  $\dim A = n$ ,  $A \subseteq [W]_{i/2^m}$ , and  $\kappa([W]_{i/2^m}, A) \leq M$ . Since  $A$  is closed  $\text{supp } W_{m,i} = \text{cl } [W]_{i/2^m}$  can be covered by the same number of translates of  $A$  as  $[W]_{i/2^m}$ , i.e.  $\kappa(\text{supp } W_{m,i}, A) = \kappa([W]_{i/2^m}, A)$ . Using Lemma 2.7 we thus obtain

$$\kappa(\text{supp } W_{m,i}, \frac{1}{2}A) \leq C_1(n)\kappa(\text{supp } W_{m,i}, A) \leq C_1(n)M .$$

Moreover,  $W_{m,i} = 1/2^m$  on  $A$  and  $W_{m,i} \leq 1/2^m$  on  $\mathbb{R}^n$ . By (iii)  $W_{m,i} \in \mathcal{W}(n, C)$  for  $C = C_1(n)^4 M$ , independently of  $m$  and  $i$ . By (ii)  $W_m \in \mathcal{W}(n, C)$  for every  $m$ , and thus (i) yields the desired result.

The remaining case (v) is handled as follows: We can assume that  $W \neq 0$ , otherwise there is nothing to do. By our assumptions there are  $M > 0$  and  $0 < t_1 < t_0$  such that  $\alpha([W]_t) \leq M$  for  $t$  in  $(0, t_1] \cup [t_0, \infty)$  and  $[W]_t \neq \emptyset$  for  $t$  in  $(0, t_1]$ . Consider  $\gamma([W]_t)$  as a function of  $t$  sending  $(0, \infty)$  into  $\{0, 1, 2, \dots, n\}$  ( $\gamma$  is given in Definition 2.5). We can choose  $0 < t_3 < t_2 \leq t_1$  with  $\gamma([W]_{t_2}) = \gamma([W]_{t_3})$ . For  $x$  in  $\mathbb{R}^n$  put  $W_1(x) := \min\{t_3, W(x)\}$  and  $W_2(x) := \min\{t_0 - t_3, W(x) - W_1(x)\}$ . Also put  $W_3 := W - W_1 - W_2$ . Then  $W_1 \leq t_3$ ,  $W_2 \leq t_0 - t_3$ , and  $W_i \geq 0$  for  $i = 1, 2, 3$ . Moreover, we have

$$[W_1]_t = \begin{cases} [W]_t & 0 \leq t \leq t_3 \\ \emptyset & t_3 < t \end{cases}$$

$$[W_2]_t = \begin{cases} [W]_{t+t_3} & 0 \leq t \leq t_0 - t_3 \\ \emptyset & t_0 - t_3 < t \end{cases}$$

$$[W_3]_t = [W]_{t+t_0} .$$

From (iv) it follows that  $W_1, W_3 \in \mathcal{W}(n, C)$  for some  $C \geq 0$ . Since  $[W]_{t_2} \neq \emptyset$  and  $\alpha([W]_{t_2}) < \infty$  there is  $A$  in  $\mathcal{C}(n)$  with  $\dim A = n$ ,  $A \subseteq [W]_{t_2}$  and  $\kappa([W]_{t_2}, A) < \infty$ . By Lemma 2.6 also  $\kappa([W]_{t_3}, A) < \infty$ , and by Lemma 2.7a)  $\kappa([W]_{t_3}, \frac{1}{2}A) < \infty$ . Hence the closedness of  $A$  and  $\text{supp } W_2 \subseteq \text{cl } [W]_{t_3}$  imply that  $\kappa(\text{supp } W_2, \frac{1}{2}A) < \infty$ . Also we have  $W_2 \geq t_2 - t_3$  on  $A$  and  $W_2 \leq t_0 - t_3$  on  $\mathbb{R}^n$ . Now (iii) implies that  $W_2 \in \mathcal{W}(n, C)$  for some  $C$ , and by (ii) the same holds for  $W = W_1 + W_2 + W_3$ . This finishes the proof of (v).

## References

- [1] Ackermann, N.: On a periodic Schrödinger equation with nonlocal superlinear part. *Math. Z.* **248**, 423–443 (2004)
- [2] Buffoni, B., Jeanjean, L., and Stuart, C.A.: Existence of a nontrivial solution to a strongly indefinite semilinear equation. *Proc. Amer. Math. Soc.* **119**, 179–186 (1993)
- [3] Grünbaum, B.: Convex polytopes. With the cooperation of Victor Klee, M. A. Perles and G. C. Shephard. *Pure and Applied Mathematics*, Vol. 16, Interscience Publishers John Wiley & Sons, Inc., New York 1967

- [4] Mattner, L.: Strict definiteness of integrals via complete monotonicity of derivatives. *Trans. Amer. Math. Soc.* **349**, 3321–3342 (1997)
- [5] Sasvári, Z.: Positive definite and definitizable functions, vol. 2 of *Mathematical Topics*. Berlin: Akademie Verlag 1994
- [6] Stewart, J.: Positive definite functions and generalizations, an historical survey. *Rocky Mountain J. Math.* **6**, 409–434 (1976)
- [7] Zong, C.: *Strange phenomena in convex and discrete geometry*. Universitext, New York: Springer-Verlag 1996

Nils Ackermann  
Mathematisches Institut  
Universität Giessen  
Arndtstr. 2  
D-35392 Giessen  
email: [nils.ackermann@math.uni-giessen.de](mailto:nils.ackermann@math.uni-giessen.de)  
web: <http://www.math.uni-giessen.de/nils.ackermann>